## RHEUMATOLOGY ENROLLMENT FORM

	BAPTIS	DATE:	 □ PAT	SHIP TO:			
					DOB □	MALE DFEMALE	
PATIENT INFO					STATE Z	'IP	
E E							
	HOME TELEPHONE	JRANCE CARD (FRONT & BACK)	MORILE PHONE		SS#		
DIAGNOSIS							
	RIOR FAILED Medication Medication						
<u>≅</u> M	EDICATION(S) Length of	f Treatmentto		Length of Tre	atment to		
Ĺ	Reason fo	or Discontinuing DIRECTIONS	QUANTITY	Reason for Di  ORENCIA®		QUANTITY	
_	162mg/0.9ml PFS	□ (wt <100kg): Inject 162mg SC every other wee		□ 250mg/15ml Vial	DIRECTIONS Starter:  Initial: Infuse mg		
	1102111g/0.21111113	$\square$ (wt >100kg): Inject 162mg SC every week	☐ 4 PFS/Pen	☐ 125mg/ml PFS	IV, then inject 125mg SC within 24 hrs	☐ 4 PFS / Pens	
		□mg/kg SC every <u>other</u> week	Refills	☐ 125mg/ml ClickJect™ Po	Maintenance. La mject 125mg 5c once a week		
	l CIMZIA®	DIRECTIONS	QUANTITY	■ OTEZLA®	☐ 0ther DIRECTIONS	Refills OUANTITY	
St	arter:	Starter Directions:	□ 1 PFS Kit (6x200 mg PFS)	☐ Starter/Titration Pack	(Use Otezla START form for bridge dosage)	☐ Starter Pack	
	l Starter Kit 200mg PFS l 200mg/ml Vial & Supplies	□ Inject 400mg SC at weeks 0, 2, and 4 □ Other:	(6x200 mg PF3)		Starter: Take as directed on Starter Pack	□30	
		I .	□ 4 Vials	□30mg Tablet	Maintenance Treatment (30mg)	60	
	aintenance l 200mg/ml PFS	Maintenance Directions: ☐ Inject 200mg SC every <u>other</u> week	☐ 6 Vials		☐ Take 1 tablet by mouth TWICE a day ☐ Take 1 tablet by mouth ONCE a day	Refills	
	l 200mg/ml Vial & Supplies	☐ Inject 400mg SC every 4 weeks	□Vials		☐ Other:		
	COSENTYX®	Other:	Refills QUANTITY	■ REMICADE®	DIRECTIONS	QUANTITY	
	COSENTIA	Starter: ☐ Ankylosing Spondylitis or Psoriatic		□ 100mg/20ml Vial	Starter: Administermg kg at 0,2, and 6 weeks, then maintenance dosing	□Vials	
_	]150mg/ml PFS	Arthritis: Inject 150mg (1 pens / PFS) SC weekly at weeks, 0,1,2,3 and 4, then maintenance dosing	☐ 5 Pen / PFS		Maintenance:	Refills	
NO	1 iovilig/iiii PF3	at weeks, 0, 1, 2, 3 and 4, then maintenance dosing	□ 10 Pens / PFS		☐ Administermg/kg every weeks		
INFORMATION	] 150mg/ml Pen	Starter: ☐ Psoriatic Arthritis with Coexistent Plaque Psoriasis: Inject 300mg (2 pens/PFS) SC	Dofile		Other:		
۷ Σ		weekly at weeks 0,1,2,3, and 4, then maintenance	Refills	■ RITUXAN® □ 100mg/10ml Vial	DIRECTIONS Starter:	QUANTITY  Vials	
DR.		• 1111		- □ 500mg/50ml Vial	Administer 1000mg IV initially and in 2 weeks	viuis	
Ĕ		Maintenance: ☐ Ankylosing Spondylitis or Psoriatic Arthritis: Inject 150mg SC every 4 weeks	□1 Pens / PFS		Maintenance:	•	
		□ Other:	☐ 2 Pens / PFS		Administer 1000 mg IV every weeks	Refills	
Z		Maintenance: □Psoriatic Arthritis with Coexisten		■ SIMPONI®	☐ Other DIRECTIONS	QUANTITY	
$\stackrel{P}{=}$		Plaque Psoriasis : Inject 300mg SC every 4 weeks.	Refills		t° (Pen) ☐ Inject 50mg SC once a month		
CRIPTION	_	□ Other:		□50mlg/0.5ml PFS	□ Other:	Refills	
S.	ENBREL®	DIRECTIONS	QUANTITY	■ SIMPONI® ARIA®	Startor: Infuso ma (2ma/ka) IV at	QUANTITY  Uials	
	l 50mg/ml SureClick Pen® l 50mg/ml PFS	□ Inject 50mg SC once a week □ Inject 50mg SC twice a week	□4 □8	□ Johng/∓illi viai	weeks 0 and 4, then maintenance dosing.	LIviais	
<u> </u>	125mg/0.5ml PFS	☐ Inject 25mg SC twice a week			Maintenance:	Refills	
	125mg/0.5ml Vial	□0.8mg/kg SC every week	Refills		☐ Infusemg (2mg/kg)IV every 8 weeks. ☐ Other:		
		□ Other:		■ STELARA®	DIRECTIONS	QUANTITY	
	HUMIRA®	DIRECTIONS	QUANTITY	☐ 45mg/0.5ml PFS	Starter: ☐ Inject 1 PFS SC on Day 1	. □ 1	
	l 40mg/0.8ml Pen l 40mg/0.8ml PFS	☐ Inject 40mg SC every <u>other</u> week ☐ Inject 40mg SC once a week	□ 2 □ 4	□90mg/1ml PFS	Maintenance: ☐ Inject 1 PFS 4 weeks after start of treatment, then every 12 weeks thereafter	t	
	r romg, o.omr r 5	□ Other	Refills		Other:	Refills	
	HUMIRA® FOR UVEITIS		QUANTITY	■XELJANZ®	DIRECTIONS	QUANTITY	
	veitis Starter ] Psoriasis Starter Pen Kit	Uveitis Induction Dose:  Inject two 40mg Pens SC on day 1, then one	☐ 1 Starter Kit	□5mg Tablet	☐ Take 5mg PO twice daily	☐ 60   Refills	
Ш,	1 L20119212 2191161 L611 KIT	40mg Pen on day 8, then one 40mg Pen every		■ XELJANZ XR®	DIRECTIONS	QUANTITY	
		other week		☐ 11mg Tablet GOUT AGENTS	☐ Take 1 tablet PO once daily  DIRECTIONS	☐ 30   Refills	
Ü	veitis Maintenance Dose:	Uveitis Maintenance Directions:	☐ 2 Pens/PFS	☐ KRYSTEXXA®	☐ Administer 8mg via iv infusion over	□2 vials	
	40mg/0.8ml Pen	☐ Inject one 40mg dose SC every other week	D	□8mg/ml Vial	2 hours every 2 weeks:	Refills	
	140mg/0.8ml PFS	Other:	Refills	n Office to must delicit	Other:	tion tuning :	
ראו	ECTION TRAINING Patient has received injection training Physician Office to provide injection training Pharmacy to provide injection training						
z	Prescriber's Name:	rescriber's Name: Contact Person:					
SER 10	Telephone:	hone: Fax: Email:					
RIE	Office Address:				City:State:	Zip:	
SC	NPI # :	DEA # :	TAX II	Medicaid Provider # :			
PRESCRIBER NFORMATION		Office Address:					
أا	PRESCRIBER'S SIGNATUR	RE (DATE)	(DATE) *IF BRAND DRUGS ARE PREFERRED, HANDWRITE "BRAND MEDICALLY NECESSARY" ABOVE				
I authorize the Pharmacy noted above and its representatives to act as an agent to initiate and execute the insurance prior authorization process.						Cept, LP All rights Reserved	